

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**THE ADSORPTION OF ADSORBABLE ORGANIC HALIDE
ONTO BIOLOGICAL SOLIDS**

A thesis submitted in partial fulfilment
of the requirements for the degree
of

MASTER OF TECHNOLOGY

in

ENVIRONMENTAL ENGINEERING

by

DANIEL JAMES GAPES

Department of Process and Environmental Technology
Massey University

March 1994

ABSTRACT

Chlorinated organic molecules may be removed from wastewaters by adsorption onto the biomass present in biological treatment systems. This study assessed the adsorption onto biological solids of Adsorbable Organic Halide (AOX) from two New Zealand kraft pulp and paper mill bleach plant wastewaters.

Batch adsorption studies were carried out to determine the characteristics of this adsorption process, with activated sludge as an adsorbent. Different molecular weight AOX fractions from two bleach plants' wastewaters were studied. The first wastewater was a combined C and E stage effluent from the CEH bleach sequence previously used at the Tasman Pulp and Paper Co. Ltd, Kawerau. The second was a combined D and E₀ stage effluent from the OODE₀D bleach sequence used at New Zealand Forest Products Kinleith Mill, Tokoroa.

For each mill's wastewater, the adsorption isotherms were characterised for four different molecular weight fractions, and the unfractionated wastewater. Adsorption isotherm models used to fit the data for each of the fractions were the Freundlich, Langmuir and Irreversible isotherms. No single model was able to successfully describe the adsorption characteristics for all of the fractions analysed, indicating significant differences in the adsorption processes occurring in the various fractions.

Analysis of the adsorption of the different molecular weight fractions demonstrated that the adsorption affinity of the AOX increased with increasing molecular weight. It appeared that molecules with higher chlorination levels were more effectively adsorbed onto the biomass. Competition for adsorption sites on the biomass by the different molecular weight fractions was a significant factor in the adsorption of the unfractionated wastewater.

Analysis of the adsorption of AOX from the wastewaters at concentrations typical of those expected in different biological treatment systems showed that:

- treatment systems with low biomass concentrations, such as aerated lagoons, would not be expected to remove significant amounts of AOX by adsorption.

- treatment systems utilising higher biomass levels have the potential to remove significant amounts of AOX by adsorption. Conventional activated sludge systems could remove 15-20%, and oxygen activated sludge systems 25-50% of the AOX entering these systems.

The impact of the modernisation of bleach sequences on the adsorption of AOX, by the introduction of oxygen delignification and chlorine dioxide substitution, was assessed. This was carried out by comparison of the adsorption characteristics of the two mills' wastewaters. The Tasman mill's bleach sequence was used as an example of older, conventional bleaching, and the Kinleith mill's sequence an example of modernised bleaching. Significant differences were found in the adsorptive behaviour of the molecular weight fractions, and that of the unfractionated wastewaters, between the two.

From the analysis, adsorption onto biomass in a biological treatment system is predicted to alter the molecular weight distribution of the AOX in wastewaters. The conventional sequence's wastewater should show a decrease in the proportion of high molecular weight AOX due to this adsorptive removal, an effect not as significant in the adsorption of AOX from a modern bleach plant wastewater. Modern bleach sequence wastewaters are expected to reveal a decrease in the proportion of the low molecular weight material, an effect not as notable for the conventional sequence's AOX.

Modernisation was found to decrease the adsorptive affinity of the wastewater, suggested to be due to the lower chlorination levels and average molecular weights of the molecules in the wastewater. It was concluded that the modernisation of bleach sequences has reduced the significance of adsorption onto biological solids as an AOX removal mechanism.

ACKNOWLEDGEMENTS

I am grateful for the support of many people in the completion of this thesis and would like to pay tribute to the following:

Firstly, my supervisors Drs Rao Bhamidimarri and Bob Chong of Massey University, and Dr Paul McFarlane of New Zealand Forest Research Institute. Their assistance during the experimental work and writing of this thesis has been invaluable.

Thanks to PAPRO New Zealand for their financial assistance.

Thanks to all the friends, flatmates, office mates and fellow postgrads who helped me keep some form of sanity throughout the course of the work.

Thanks to my parents, John and Jan, who have supported me through the whole of my life.

Thanks to Nicola, who helped me with the final preparation of the thesis, not a small task at all.

Finally, thanks be to God.

TABLE OF CONTENTS

ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
TABLE OF CONTENTS	v
LIST OF FIGURES	ix
LIST OF TABLES	x
 1. INTRODUCTION	 1
1.1. THE NEW ZEALAND PULP AND PAPER INDUSTRY	1
1.1.1. Tasman Pulp and Paper Company Ltd.....	1
1.1.2. NZFP Pulp and Paper Ltd, Kinleith Industries	2
1.2. ADSORBABLE ORGANIC HALIDE	2
1.3. OBJECTIVES OF THE WORK.....	4
2. LITERATURE REVIEW	6
2.1. GENERAL INTRODUCTION.....	6
2.2. OVERVIEW OF CHEMICAL PULPING.....	6
2.3. OVERVIEW OF PULP BLEACHING.....	7
2.3.1. Introduction.....	7
2.3.2. Lignin Structure	7
2.3.3. Bleaching Stages used in the Pulp and Paper Industry	9
2.3.3.1. Chlorine (C)	9
2.3.3.2. Alkali Extraction (E)	9
2.3.3.3. Chlorine Dioxide (D).....	12
2.3.3.4. Oxygen (O).....	12
2.3.3.5. Hydrogen Peroxide (P)	13
2.3.3.6. Ozone (Z)	13
2.4. NATURE OF BLEACHING EFFLUENTS.....	13
2.4.1. Introduction.....	13
2.4.2. High Molecular Weight Compounds	14
2.4.3. Low Molecular Weight Compounds	15
2.4.3.1. Acidic Compounds	16
2.4.3.2. Phenolic compounds	16
2.4.3.3. Neutral Compounds	16
2.4.4. Environmental Effects of Bleach Effluents	17
2.4.5. Toxic Constituents of Bleach Plant Effluents	18
2.5. REDUCTION OF CHLOROORGANICS IN BLEACH EFFLUENTS.....	19

2.5.1. Introduction.....	19
2.5.2. In-Process Reduction.....	19
2.5.3. Effluent Treatment.....	20
2.5.3.1. Gas Stripping.....	21
2.5.3.2. Precipitation.....	22
2.5.3.3. Adsorption onto Biomass.....	22
2.5.3.4. Abiotic and Biological Mineralisation.....	22
2.6. AOX REMOVALS IN ACTIVATED SLUDGE PLANTS AND AERATED LAGOONS	23
2.7. ADSORPTION OF ORGANIC POLLUTANTS.....	26
2.7.1. Adsorption Phenomena.....	26
2.7.2. Adsorption Equilibria	27
2.7.2.1. Linear Isotherm.....	28
2.7.2.2. Langmuir Isotherm.....	28
2.7.2.3. Freundlich Isotherm.....	29
2.7.2.4. Irreversible Isotherm	30
2.7.3. Mechanistic Models	30
2.7.3.1. Ion Exchange and Surface Complexation.....	30
2.7.3.2. Solvophobic Adsorption	31
2.7.4. Sorption Rate Processes	32
2.7.5. Factors Affecting Adsorption.....	33
2.7.5.1. Nature of the Adsorbate	33
2.7.5.2. Nature of the Adsorbent	33
2.7.5.3. Surface Area.....	33
2.7.5.4. Temperature.....	34
2.7.6. Adsorption of Mixed Solutes.....	34
2.7.7. Sorption of Organics onto Wastewater Solids.....	34
2.7.8. Adsorption of AOX onto Biological Solids.....	36
3. MATERIALS AND METHODS.....	40
3.1. ULTRAFILTRATION OF BLEACH PLANT SAMPLES	40
3.2. SAMPLE DEFINITION.....	41
3.3. PH ADJUSTMENT	42
3.4. DETERMINATION OF AOX CONCENTRATION.....	42
3.5. REJECTION COEFFICIENT DETERMINATION.....	42
3.6. DETERMINATION OF CONDUCTIVITY	43
3.7. COLLECTION AND STORAGE OF BIOMASS	45
3.8. MEASUREMENT OF ADSORPTION ISOTHERMS	45

3.8.1. Tasman Effluents.....	45
3.8.2. Kinleith Effluents.....	50
3.9. MODIFICATION TO MLSS PREPARATION PROCEDURE	53
3.10. ADSORPTION RATE EXPERIMENTS.....	53
3.11. BLEACH PLANT DEFINITION.....	54
3.12. DETERMINATION OF ADSORPTION ISOTHERM MODELS	54
3.12.1. Introduction.....	54
3.12.2. Langmuir Isotherm	55
3.12.3. Freundlich Isotherm	56
3.12.4. Irreversible Isotherm.....	56
4. RESULTS AND DISCUSSION.....	57
4.1. AOX DETERMINATION	57
4.2. ULTRAFILTRATION OF SAMPLES.....	58
4.2.1. Introduction.....	58
4.2.2. Ultrafiltration of the Wastewaters.....	59
4.2.3. Rejection Coefficient Analysis.....	59
4.3. COMPARISONS BETWEEN TASMAN AND KINLEITH FRACTIONS	61
4.3.1. AOX Distribution.....	61
4.3.2. Suspended Solids	63
4.4. ADSORPTION RATE EXPERIMENTS.....	64
4.5. DETERMINATION OF ISOTHERMS.....	68
4.5.1. Introduction.....	68
4.5.2. Langmuir Isotherm	77
4.5.3. Irreversible Isotherm.....	78
4.5.4. Freundlich Isotherm	80
4.5.5. Kinleith <2kD Fraction	81
4.5.6. Summary of the Adsorption Isotherms	82
4.6. COMPARISON OF ADSORPTION CHARACTERISTICS BETWEEN MILLS.....	83
4.6.1. Introduction.....	83
4.6.2. <2kD Fractions	83
4.6.3. 2-30kD Fractions.....	87
4.6.4. 30-100kD Fractions.....	89
4.6.5. >100kD Fractions	89
4.6.6. Unfractionated Samples	92

4.6.7. Overall Comparison of Adsorptive Behaviour Between the Two Mill's Effluents	94
4.7. COMPARISON OF ADSORPTION CHARACTERISTICS: WITHIN MILL SAMPLES.....	95
4.7.1. Tasman.....	95
4.7.2. Kinleith Samples	99
4.8. IMPLICATIONS OF RESULTS IN BIOLOGICAL TREATMENT OF PULP AND PAPER WASTEWATERS	101
5. CONCLUSIONS AND RECOMMENDATIONS	109
5.1. CONCLUSIONS	109
5.2. RECOMMENDATIONS FOR FURTHER WORK	112
6. REFERENCES.....	114
APPENDIX	120

LIST OF FIGURES

Figures

	Page
1. Lignin Precursors.....	8
2. Prominent Structures in Softwood Lignin.....	8
3. Substitution and Dealkylation Reactions of Chloronium Ion	10
4. Addition Reactions of Chloronium Ion to C=C Bond.....	10
5. Addition Reactions of Chloronium Ion to other Ring Conjugated Structures.....	11
6. Displacement of Chlorine by OH ⁻ Ions, and Alkali Promoted Rearrangements.....	11
7. Phenolic Compounds Most Frequently Identified in Spent Bleaching Liquors.....	16
8. Anaerobic Removal of AOX During the Batch Digestion of Different Molecular Weight Fractions of Bleach Plant Effluent.....	23
9. Generalised Equilibrium Adsorption Isotherms.....	28
10. Schematic Representation of an Electrical Double Layer.....	31
11. Representation of Simple Cell present in Wastewater Treatment Systems.....	35
12. Standard Curve for Conductivity of Sodium Chloride Solution	44
13. AOX Distribution of Unfractionated Samples	60
14. Molecular Weight Distribution of Wastewater Samples	60
15. Proportional Distribution of Total AOX and TSS Present in Each Fraction	60
16. Adsorption - Time Plots for Tasman <2kD and >100kD Fractions.....	65
17. Modelled Adsorption Rates for Tasman <2 and >100kD Fractions	65
18. Percentage Adsorption-Time Plots for Tasman <2kD and >100kD Fractions	65
19. Adsorption of Tasman <2kD Fraction	69
20. Adsorption of Tasman 2-30kD Fraction.....	69
21. Adsorption of Tasman 30-100kD Fraction.....	69
22. Adsorption of Tasman >100kD Fraction	70
23. Adsorption of Tasman Unfractionated Sample.....	70
24. Adsorption of Kinleith <2kD Fraction	70
25. Adsorption of Kinleith 2-30kD Fraction.....	71
26. Adsorption of Kinleith 30-100kD Fraction.....	71

27.	Adsorption of Kinleith >100kD Fraction	71
28.	Adsorption of Kinleith Unfractionated Sample	72
29.	Adsorption Isotherm, Tasman <2kD Fraction	73
30.	Adsorption Isotherm, Tasman 2-30kD Fraction	73
31.	Adsorption Isotherm, Tasman 30-100kD Fraction	73
32.	Adsorption Isotherm, Tasman >100kD Fraction	74
33.	Adsorption Isotherm, Tasman Unfractionated Sample	74
33b.	Adsorption Isotherm, Tasman Unfractionated Sample using Model Co Value	74
34.	Adsorption Isotherm, Kinleith <2kD Fraction	75
35.	Adsorption Isotherm, Kinleith 2-30kD Fraction	75
36.	Adsorption Isotherm, Kinleith 30-100kD Fraction	75
36b.	Adsorption Isotherm, Kinleith 30-100kD Fraction using model Co.	76
37.	Adsorption Isotherm, Kinleith >100kD Fraction	76
38.	Adsorption Isotherm, Kinleith Unfractionated Sample	76
39.	Adsorption of <2kD Fractions	85
40.	Relative Adsorption of <2kD Fractions	85
41.	Adsorption Isotherm, <2kD Fractions	85
42.	Adsorption of 2-30kD Fractions	88
43.	Relative Adsorption of 2-30kD Fractions	88
44.	Adsorption Isotherm, 2-30kD Fractions	88
45.	Adsorption of 30-100kD Fractions	90
46.	Relative Adsorption of 30-100kD Fractions	90
47.	Adsorption Isotherm, 30-100kD Fractions	90
48.	Adsorption of >100kD Fractions	91
49.	Relative Adsorption of >100kD Fractions	91
50.	Adsorption Isotherm, >100kD Fractions	91
51.	Adsorption of Unfractionated Samples	93
52.	Relative Adsorption of Unfractionated Samples	93
53.	Adsorption Isotherm, Unfractionated Samples	93
54.	Adsorption Isotherm, Tasman Samples	96
55.	Relative Adsorption of Tasman Samples	96
56.	Adsorption of Tasman Samples	96
57.	Adsorption Isotherm, Kinleith Samples	100
58.	Relative Adsorption of Kinleith Samples	100
59.	Adsorption of Kinleith Samples	100
60.	Tasman AOX Loadings at Different Biomass Concentrations	103

LIST OF TABLES

Tables

	Page
I. Pulp and Paper Mills in New Zealand	2
II. Substances Dissolved during Bleaching of Kraft Pulps	14
III. Some Compounds Identified in C and E Stage Wastewaters.....	15
IV. Removal of AOX from Biological Treatment Systems	24
V. Freundlich Parameters for Adsorption of Chloroorganics onto Biological Solids.....	37
VI. Samples from Ultrafiltration Procedure	41
VII. Dilution and Phosphate Buffer Addition of Tasman AOX Samples	48
VIII. Dilution and Phosphate Buffer Addition of Kinleith AOX Samples	51
IX. Apparent Rejection Coefficients (R) for Wastewater Fractions	59
X. AOX Distribution in Kinleith Mill Wastewaters.....	63
XI. Calculated Rate Parameters for Tasman Wastewater Fractions	66
XII. Results of Langmuir Adsorption Isotherm Analysis	77
XIII. Results of Irreversible Adsorption Isotherm Analysis	78
XIV. Results of Freundlich Adsorption Isotherm Analysis.....	80
XV. Adsorption Isotherm model for bleach plant AOX.....	82
XVI. Average Ratios of TOX/TOC of Kraft Mill Wastewater	92
XVII. Characteristic Parameters of Three Treatment Systems	102
XVIII. Percent of AOX in Each Fraction that could Potentially be Adsorbed in Different Treatment Systems	105
XIX. Adsorption Isotherms for Bleach Plant AOX.....	109